# INTERNAL WAVE MOORING

Murray Levine541-737-3047 levine@oce.orst.edufax: 541-737-2064 Timothy Boyd541-737-4035tboyd@oce.orst.edufax: 541-737-2064

104 Ocean Admin Bldg Oregon State University Corvallis, OR 97331

Award #: N00014-95-1-0534

### LONG-TERM GOALS

The overall goal of this PRIMER program is to assess the feasibility of the operation of a synthetic aperture sonar on the continental shelf. To evaluate the performance of this sonar system, knowledge of the fluctuations of oceanic sound speed, primarily due to internal waves, is needed.

### SCIENTIFIC OBJECTIVES

The specific objectives of this project are:

- to describe the internal wave field and associated sound speed fluctuations on the continental shelf--both statistically and by events, and
- to understand the processes involved in the transition of the internal wave field from the deep ocean to the continental slope and shelf.

## **APPROACH**

We deployed a densely-instrumented sub-surface mooring over the continental shelf to record time series of velocity and temperature at vertical and temporal scales appropriate to resolve the internal wave field. A crude estimate of the horizontal correlation of the wave field was made using a reduced set of instruments on two surface buoys within 300 m of the main mooring. The moorings were operating during the PRIMER acoustic observations made by K. Williams, F. Henyey (UW/APL), and S. Stanic (URL). The field work was conducted cooperatively with investigators involved in the ONR Coastal Mixing & Optics program--especially T. Dickey (UCSB), and S. Lentz, A. Plueddemann, S. Andersen (WHOI).

## WORK COMPLETED

The moorings were deployed on the shelf in 70 m of water on 9 July 1996 and recovered about 3 months later on 26 Sept 1996. Almost all of the instruments functioned properly, recording data every 2 minutes. A data report has been published (Boyd, Levine, and Gard, 1997) and a CD containing converted data has been released. Preliminary results were presented at several workshops and at The Oceanography Society meeting in Seattle, 1997.

maintaining the data needed, and c including suggestions for reducing	lection of information is estimated to ompleting and reviewing the collect this burden, to Washington Headqu uld be aware that notwithstanding an DMB control number.	ion of information. Send comment arters Services, Directorate for Info	s regarding this burden estimate ormation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE 30 SEP 1997		2. REPORT TYPE		3. DATES COVERED <b>00-00-1997 to 00-00-1997</b>		
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER		
Internal Wave Mooring				5b. GRANT NUMBER		
				5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)				5d. PROJECT NUMBER		
				5e. TASK NUMBER		
				5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  Oregon State University, College of Oceanic and Atmospheric Sciences, 104 Ocean Admin. Bldg., Corvallis, OR, 97331				8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)		
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAIL Approved for publ	ABILITY STATEMENT ic release; distributi	ion unlimited				
13. SUPPLEMENTARY NO	OTES					
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>	Same as Report (SAR)	3		

**Report Documentation Page** 


Form Approved OMB No. 0704-0188

# **RESULTS**


The 3-month time series captured a variety of different phenomena that span a range of space and time scales. Some examples can be seen in the temperature time series shown here.

Two major storm systems, which were downgraded hurricanes, passed near the mooring site. These storms resulted in significant mixing events (Figure 1). The temperature at 11 m typically cooled during the storms as colder water was entrained from below. During Edouard the temperature as deep as 54 m warmed apparently due to vertical mixing. The change in potential energy due to the passage of Edouard was calculated and compared with expectations from simple bulk models.

Tidal oscillations are prevalent in the temperature at 11 m and are primarily due to the vertical displacement by the internal tide. After the mixing of the upper layer by Bertha and Edouard, the tidal oscillations are much reduced; the internal tide may still exist, but the vertical temperature gradient is much weaker.



*Figure 1.* Temperature time series recorded at 11 and 54 m. The arrival times of tropical storms Bertha and Edouard are indicated.



*Figure 2.* A 12-hour time series of temperature recorded at 4 depths during the passage of a packet of high-frequency internal waves. The packet consists of a series of sharp downward displacement "spikes" which correspond to sharp temperature increases.

High-frequency nonlinear wave packets occur throughout the time series. These packets are generated by the interaction of the barotropic tide with the shelf break. Packets are usually separated by multiples of a tidal period (about 12 hours). A typical example of a wave packet is shown in Figure 2. The wave period is about 12 minutes with vertical displacements reaching 10 m. More will be learned when the complete set of temperature time series (every 3 to 6 m in the vertical) are analyzed in conjunction with the velocity observations (every 4 m).

## **IMPACT / APPLICATIONS**

The measured oceanic variability will be compared with observed acoustic observations and used as input to acoustic models in order to assess the potential performance of synthetic aperture sonar systems on the continental shelf. We will also use the detailed analysis of the internal wave field to provide a framework for creating a climatology of internal waves on the shelf and slope.

### **TRANSITIONS**

The observed variability of the internal wave field will be used as input to the acoustic modeling effort, which ultimately will assess the feasibility of a SAS system in this environment. Defining the internal wave field is also important to those investigators who measured turbulent mixing and optical fluctuations at the same time under related programs.

## RELATED PROJECTS

The data from the moorings are being shared with many investigators involved in the Coastal Mixing & Optics program who made complementary observations during summer 1996, including T. Dickey (UCSB), R. Zaneveld, J. Barth, M. Kosro (OSU), S. Lentz, A. Plueddemann, S. Andersen (WHOI). We look forward to participating in joint analysis where appropriate combining our internal wave observations with the optical and turbulence data.

### REFERENCES

Boyd, T., M.D. Levine, and S.R. Gard, Mooring observations from the Mid-Atlantic Bight, July-September 1996, Ref. 97-2, Data report 164, Oregon State University, 1997.